

# Perturbed Inertial Krasnoselskii-Mann Iterations and its application to image inpainting

Daniel Cortild

Faculty of Science and Engineering  
University of Groningen

Optimization at Bernoulli Institute  
Groningen (Netherlands), June 18-20, 2023



university of  
groningen

faculty of science  
and engineering

mathematics and applied  
mathematics

# Optimisation and Fixed Point

Let  $f: \mathcal{H} \rightarrow \mathbb{R}$  be differentiable.

Consider the problem: Find  $\hat{x}$  such that

$$f(\hat{x}) = \min f(x)$$

# Optimisation and Fixed Point

Let  $f: \mathcal{H} \rightarrow \mathbb{R}$  be differentiable.

Consider the problem: Find  $\hat{x}$  such that

$$f(\hat{x}) = \min f(x)$$

Equivalent problem: Find  $\hat{x}$  such that

$$\nabla f(\hat{x}) = 0$$

# Optimisation and Fixed Point

Let  $f: \mathcal{H} \rightarrow \mathbb{R}$  be differentiable.

Consider the problem: Find  $\hat{x}$  such that

$$f(\hat{x}) = \min f(x)$$

Equivalent problem: Find  $\hat{x}$  such that

$$\nabla f(\hat{x}) = 0$$

Other equivalent problem: Find  $\hat{x}$  such that

$$T\hat{x} = \hat{x}, \quad \text{where } T = I - \nabla f$$

# Optimisation and Fixed Point

Let  $f, g: \mathcal{H} \rightarrow \mathbb{R}$ . Suppose that  $f$  is differentiable.

Consider the problem: Find  $\hat{x}$  such that

$$(f + g)(\hat{x}) = \min f(x) + g(x)$$

# Optimisation and Fixed Point

Let  $f, g: \mathcal{H} \rightarrow \mathbb{R}$ . Suppose that  $f$  is differentiable.

Consider the problem: Find  $\hat{x}$  such that

$$(f + g)(\hat{x}) = \min f(x) + g(x)$$

Equivalent problem: Find  $\hat{x}$  such that

$$\begin{aligned} 0 &\in \partial(f + g)(\hat{x}) = \nabla f(\hat{x}) + \partial g(\hat{x}) \\ \iff \hat{x} &= (I - \partial g)^{-1}(I + \nabla f)(\hat{x}) \end{aligned}$$

# Optimisation and Fixed Point

Let  $f, g: \mathcal{H} \rightarrow \mathbb{R}$ . Suppose that  $f$  is differentiable.

Consider the problem: Find  $\hat{x}$  such that

$$(f + g)(\hat{x}) = \min f(x) + g(x)$$

Equivalent problem: Find  $\hat{x}$  such that

$$\begin{aligned} 0 &\in \partial(f + g)(\hat{x}) = \nabla f(\hat{x}) + \partial g(\hat{x}) \\ \iff \hat{x} &= (I - \partial g)^{-1}(I + \nabla f)(\hat{x}) \end{aligned}$$

Other equivalent problem: Find  $\hat{x}$  such that

$$T\hat{x} = \hat{x}, \quad \text{where } T = (I - \partial g)^{-1}(I + \nabla f)$$

# Fixed Point Iterative Schemes

## Problem

For  $T: \mathcal{H} \rightarrow \mathcal{H}$ , find  $\hat{x} \in \mathcal{H}$  such that  $T\hat{x} = \hat{x}$ .



# Fixed Point Iterative Schemes

## Problem

For  $T: \mathcal{H} \rightarrow \mathcal{H}$ , find  $\hat{x} \in \mathcal{H}$  such that  $T\hat{x} = \hat{x}$ .

Picard Iterations:

$$x_{k+1} = Tx_k$$

# Fixed Point Iterative Schemes

## Problem

For  $T: \mathcal{H} \rightarrow \mathcal{H}$ , find  $\hat{x} \in \mathcal{H}$  such that  $T\hat{x} = \hat{x}$ .

Picard Iterations:

$$x_{k+1} = Tx_k$$

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k Tx_k$$

# Fixed Point Iterative Schemes

## Problem

For  $T: \mathcal{H} \rightarrow \mathcal{H}$ , find  $\hat{x} \in \mathcal{H}$  such that  $T\hat{x} = \hat{x}$ .

Picard Iterations:

$$x_{k+1} = Tx_k$$

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k Tx_k$$

Advantages of KM iterations over Picard iterations:

# Fixed Point Iterative Schemes

## Problem

For  $T: \mathcal{H} \rightarrow \mathcal{H}$ , find  $\hat{x} \in \mathcal{H}$  such that  $T\hat{x} = \hat{x}$ .

Picard Iterations:

$$x_{k+1} = Tx_k$$

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k Tx_k$$

Advantages of KM iterations over Picard iterations:

- Allows for a broader class of operators.

# Fixed Point Iterative Schemes

## Problem

For  $T: \mathcal{H} \rightarrow \mathcal{H}$ , find  $\hat{x} \in \mathcal{H}$  such that  $T\hat{x} = \hat{x}$ .

Picard Iterations:

$$x_{k+1} = Tx_k$$

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k Tx_k$$

Advantages of KM iterations over Picard iterations:

- Allows for a broader class of operators.
- Includes more diverse algorithms.

# Acceleration Methods

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

# Acceleration Methods

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

Heavy-Ball Acceleration:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T x_k \end{cases}$$

# Acceleration Methods

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

Heavy-Ball Acceleration:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T x_k \end{cases}$$

Nesterov Acceleration:

$$\begin{cases} z_k &= x_k + \beta_k(x_k - x_{k-1}) \\ x_{k+1} &= (1 - \lambda_k)z_k + \lambda_k T z_k \end{cases}$$



# Acceleration Methods

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

General Inertial Scheme:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T z_k \end{cases}$$

# Acceleration Methods

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

General Inertial Scheme:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T z_k \end{cases}$$

Special cases:

# Acceleration Methods

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

General Inertial Scheme:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T z_k \end{cases}$$

Special cases:

- When  $\alpha_k = \beta_k = 0$ , we get no acceleration

# Acceleration Methods

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

General Inertial Scheme:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T z_k \end{cases}$$

Special cases:

- When  $\alpha_k = \beta_k = 0$ , we get no acceleration
- When  $\alpha_k = \beta_k$ , we get Nesterov acceleration

# Acceleration Methods

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

General Inertial Scheme:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T z_k \end{cases}$$

Special cases:

- When  $\alpha_k = \beta_k = 0$ , we get no acceleration
- When  $\alpha_k = \beta_k$ , we get Nesterov acceleration
- When  $\beta_k = 0$ , we get Heavy-Ball acceleration

# Acceleration Methods

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

General Inertial Scheme:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T z_k \end{cases}$$

Special cases:

- When  $\alpha_k = \beta_k = 0$ , we get no acceleration
- When  $\alpha_k = \beta_k$ , we get Nesterov acceleration
- When  $\beta_k = 0$ , we get Heavy-Ball acceleration
- When  $\alpha_k = 0$ , we get reflected acceleration

# Adding Perturbations

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

# Adding Perturbations

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

Perturbed Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k + \varepsilon_k$$



# Adding Perturbations

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

Perturbed Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k + \varepsilon_k$$

Let  $T_k = T + E_k$  where  $E_k \rightarrow 0$ .

# Adding Perturbations

Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k$$

Perturbed Krasnoselskii-Mann Iterations:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T x_k + \varepsilon_k$$

Let  $T_k = T + E_k$  where  $E_k \rightarrow 0$ . KM iterations with the operators  $T_k$  replacing  $T$  may be written as

$$\begin{aligned} x_{k+1} &= (1 - \lambda_k)x_k + \lambda_k T_k x_k \\ &= (1 - \lambda_k)x_k + \lambda_k T x_k + \lambda_k E_k x_k \\ &= (1 - \lambda_k)x_k + \lambda_k T x_k + \varepsilon_k, \end{aligned}$$

with  $\varepsilon_k = \lambda_k E_k x_k$ .

# Convergence Theorems

Perturbed General Inertial KM Iterations:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) + \varepsilon_k \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) + \rho_k \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k Tz_k + \theta_k \end{cases}$$

# Convergence Theorems

Perturbed General Inertial KM Iterations:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) + \varepsilon_k \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) + \rho_k \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k Tz_k + \theta_k \end{cases}$$

## Theorem (Weak Convergence)

Let  $T: \mathcal{H} \rightarrow \mathcal{H}$  be nonexpansive such that  $F := \text{Fix}(T) \neq \emptyset$ . Under mild conditions,  $(x_k)$ ,  $(y_k)$  and  $(z_k)$  converge weakly to a same point in  $F$ .

# Convergence Theorems

Perturbed General Inertial KM Iterations:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) + \varepsilon_k \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) + \rho_k \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k Tz_k + \theta_k \end{cases}$$

## Theorem (Weak Convergence)

Let  $T: \mathcal{H} \rightarrow \mathcal{H}$  be nonexpansive such that  $F := \text{Fix}(T) \neq \emptyset$ . *Under mild conditions*,  $(x_k)$ ,  $(y_k)$  and  $(z_k)$  converge weakly to a same point in  $F$ .

## Theorem (Strong Convergence)

Let  $T: \mathcal{H} \rightarrow \mathcal{H}$  be contractive such that  $\text{Fix}(T) = \{p^*\}$ . *Under mild conditions*,  $(x_k)$  converges strongly to  $p^*$ .

# Extensions

The previous theorems may be extended by

# Extensions

The previous theorems may be extended by

- ① replacing “nonexpansive” and “contractive” by “quasi-nonexpansive” ( $K = 1$ ) and “quasi-contractive” ( $K < 1$ ):

$$\|Tx - p\| \leq K\|x - p\|, \quad \forall x \in \mathcal{H}, p \in \text{Fix}(T)$$

# Extensions

The previous theorems may be extended by

- ① replacing “nonexpansive” and “contractive” by “quasi-nonexpansive” ( $K = 1$ ) and “quasi-contractive” ( $K < 1$ ):

$$\|Tx - p\| \leq K\|x - p\|, \quad \forall x \in \mathcal{H}, p \in \text{Fix}(T)$$

- ② taking a family of operators  $T_k: \mathcal{H} \rightarrow \mathcal{H}$  such that  $F := \bigcap_{k \geq 1} \text{Fix}(T_k) \neq \emptyset$  (Weak case) or  $\text{Fix}(T_k) = \{p^*\}$  for all  $k \geq 1$  (Strong case).



# Extensions

The previous theorems may be extended by

- ① replacing “nonexpansive” and “contractive” by “quasi-nonexpansive” ( $K = 1$ ) and “quasi-contractive” ( $K < 1$ ):

$$\|Tx - p\| \leq K\|x - p\|, \quad \forall x \in \mathcal{H}, p \in \text{Fix}(T)$$

- ② taking a family of operators  $T_k: \mathcal{H} \rightarrow \mathcal{H}$  such that  $F := \bigcap_{k \geq 1} \text{Fix}(T_k) \neq \emptyset$  (Weak case) or  $\text{Fix}(T_k) = \{p^*\}$  for all  $k \geq 1$  (Strong case).
- ③ taking a family of operators  $T_k: \mathcal{H} \rightarrow \mathcal{H}$  such that  $F := \text{Li}(\text{Fix}(T_k)) \neq \emptyset$  (Weak case) or  $\text{Fix}(T_k) = \{p_k\}$  such that  $p_k \rightarrow p^*$  (Strong case).

# Extended Convergence Theorems

Perturbed General Inertial KM Iterations:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) + \varepsilon_k \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) + \rho_k \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T_k z_k + \theta_k \end{cases}$$

# Extended Convergence Theorems

Perturbed General Inertial KM Iterations:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) + \varepsilon_k \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) + \rho_k \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T_k z_k + \theta_k \end{cases}$$

## Extended Theorem (Weak Convergence)

Let  $T_k: \mathcal{H} \rightarrow \mathcal{H}$  be quasi-nonexpansive such that  $F := \text{Li}(\text{Fix}(T_k)) \neq \emptyset$ .  
Under mild conditions,  $(x_k)$ ,  $(y_k)$  and  $(z_k)$  converge weakly to a same point in  $F$ .

# Extended Convergence Theorems

Perturbed General Inertial KM Iterations:

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) + \varepsilon_k \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) + \rho_k \\ x_{k+1} &= (1 - \lambda_k)y_k + \lambda_k T_k z_k + \theta_k \end{cases}$$

## Extended Theorem (Weak Convergence)

Let  $T_k: \mathcal{H} \rightarrow \mathcal{H}$  be quasi-nonexpansive such that  $F := \text{Li}(\text{Fix}(T_k)) \neq \emptyset$ . Under mild conditions,  $(x_k)$ ,  $(y_k)$  and  $(z_k)$  converge weakly to a same point in  $F$ .

## Extended Theorem (Strong Convergence)

Let  $T_k: \mathcal{H} \rightarrow \mathcal{H}$  be quasi-contractive such that  $\text{Fix}(T_k) = \{p_k\}$  and  $p_k \rightarrow p^*$ . Under mild conditions,  $(x_k)$  converges strongly to  $p^*$ .

# Application to Optimisation

## Problem

Let  $f, g: \mathcal{H} \rightarrow \mathbb{R} \cup \{+\infty\}$ ,  $h: \mathcal{H} \rightarrow \mathbb{R}$ , and  $L: \mathcal{H} \rightarrow \mathcal{H}$ . Find

$$\min_{x \in \mathcal{H}} f(x) + g(x) + h(Lx).$$

# Application to Optimisation

## Problem

Let  $f, g: \mathcal{H} \rightarrow \mathbb{R} \cup \{+\infty\}$ ,  $h: \mathcal{H} \rightarrow \mathbb{R}$ , and  $L: \mathcal{H} \rightarrow \mathcal{H}$ . Find

$$\min_{x \in \mathcal{H}} f(x) + g(x) + h(Lx).$$

This may be solved by the **three-operator splitting method** (Davis, Yin, 2017):

$$T_k := I - \text{prox}_{\rho_k g} + \text{prox}_{\rho_k f} \circ (2\text{prox}_{\rho_k g} - I - \rho_k L^* \circ \nabla h \circ L \circ \text{prox}_{\rho_k g}).$$

# Application to Optimisation

## Problem

Let  $f, g: \mathcal{H} \rightarrow \mathbb{R} \cup \{+\infty\}$ ,  $h: \mathcal{H} \rightarrow \mathbb{R}$ , and  $L: \mathcal{H} \rightarrow \mathcal{H}$ . Find

$$\min_{x \in \mathcal{H}} f(x) + g(x) + h(Lx).$$

This may be solved by the **three-operator splitting method** (Davis, Yin, 2017):

$$\begin{cases} y_k &= x_k + \alpha_k(x_k - x_{k-1}) + \varepsilon_k \\ z_k &= x_k + \beta_k(x_k - x_{k-1}) + \rho_k \\ z_k^g &= \text{prox}_{\rho_k g}(z_k) \\ z_k^f &= \text{prox}_{\rho_k f}(2z_k^g - z_k - \rho_k L^* \circ \nabla h \circ L(z_k^g)) \\ x_{k+1} &= y_k + \lambda_k(z_k^f - z_k^g) + \theta_k \end{cases}$$

# Image Inpainting?

Original Image





# Image Inpainting?

Original Image



Corrupt Image



# Image Inpainting?

Original Image



Corrupt Image



Recovered Image



# Image Inpainting?

Original Image



Corrupt Image



Recovered Image



**Figure:** Not obtained through described algorithm!

# Image Inpainting?

Original Image



Corrupt Image



Recovered Image



Figure: Not obtained through described algorithm!

## Mathematical Formulation

$$\min_{Z \in [0,1]^{M \times N \times 3}} \left\{ \frac{1}{2} \|\mathcal{A}Z - Z_{\text{corrupt}}\|^2 + \sigma \|Z_{(1)}\|_* + \sigma \|Z_{(2)}\|_* \right\}$$

# Visual Results

Original Image



Corrupt Image



Heavy-Ball



Perturbed Heavy-Ball



Nesterov



Perturbed Nesterov



**Figure:** Process obtained with  $\rho = 1.8$ ,  $\lambda = 1.3$ , and  $\sigma = 0.5$ .

# Convergence Plots




Figure: Process obtained with  $\rho = 1.8$ ,  $\lambda = 1.3$ , and  $\sigma = 0.5$ .

# Result Based on Algorithm

Original Image



Corrupt Image



Recovered Image



**Figure:** Obtained through perturbed inertial algorithm.

# Conclusion

Summary:



# Conclusion

## Summary:

- Perturbed Inertial KM Iterations generalise previously known algorithms.

# Conclusion

## Summary:

- Perturbed Inertial KM Iterations generalise previously known algorithms.
- They incorporate multiple types of acceleration.

# Conclusion

## Summary:

- Perturbed Inertial KM Iterations generalise previously known algorithms.
- They incorporate multiple types of acceleration.
- They account for (rounding) errors and/or inexact computations.

# Conclusion

## Summary:

- Perturbed Inertial KM Iterations generalise previously known algorithms.
- They incorporate multiple types of acceleration.
- They account for (rounding) errors and/or inexact computations.
- They also allow for operators not sharing a common fixed point.

# Conclusion

## Summary:

- Perturbed Inertial KM Iterations generalise previously known algorithms.
- They incorporate multiple types of acceleration.
- They account for (rounding) errors and/or inexact computations.
- They also allow for operators not sharing a common fixed point.
- They converge weakly and strongly under mild conditions.

# Conclusion

## Summary:

- Perturbed Inertial KM Iterations generalise previously known algorithms.
- They incorporate multiple types of acceleration.
- They account for (rounding) errors and/or inexact computations.
- They also allow for operators not sharing a common fixed point.
- They converge weakly and strongly under mild conditions.
- They include the three-operator splitting method, used for optimisation problems such as the image inpainting problem.

# Conclusion

## Summary:

- Perturbed Inertial KM Iterations generalise previously known algorithms.
- They incorporate multiple types of acceleration.
- They account for (rounding) errors and/or inexact computations.
- They also allow for operators not sharing a common fixed point.
- They converge weakly and strongly under mild conditions.
- They include the three-operator splitting method, used for optimisation problems such as the image inpainting problem.

Further possible research:

# Conclusion

## Summary:

- Perturbed Inertial KM Iterations generalise previously known algorithms.
- They incorporate multiple types of acceleration.
- They account for (rounding) errors and/or inexact computations.
- They also allow for operators not sharing a common fixed point.
- They converge weakly and strongly under mild conditions.
- They include the three-operator splitting method, used for optimisation problems such as the image inpainting problem.

## Further possible research:

- Study of the rates of convergence.



# Conclusion

## Summary:

- Perturbed Inertial KM Iterations generalise previously known algorithms.
- They incorporate multiple types of acceleration.
- They account for (rounding) errors and/or inexact computations.
- They also allow for operators not sharing a common fixed point.
- They converge weakly and strongly under mild conditions.
- They include the three-operator splitting method, used for optimisation problems such as the image inpainting problem.

## Further possible research:

- Study of the rates of convergence.
- Why do the perturbations help?

Thank you!